



Two-dimensional electron gases at a spinel/perovskite complex oxide heterointerface with electron mobilities exceeding 100,000 cm²V⁻¹s⁻¹

Chen, Yunzhong; Trier, Felix; Christensen, Dennis Valbjørn; Linderroth, Søren; Pryds, Nini

Published in:

Proceedings for the international conference: "MAMA-Trend: Trends, challenges and emergent new phenomena in multi-functional materials"

Publication date:

2013

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Chen, Y., Trier, F., Christensen, D. V., Linderroth, S., & Pryds, N. (2013). Two-dimensional electron gases at a spinel/perovskite complex oxide heterointerface with electron mobilities exceeding 100,000 cm²V⁻¹s⁻¹. In *Proceedings for the international conference: "MAMA-Trend: Trends, challenges and emergent new phenomena in multi-functional materials"* (pp. 77)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

4.5 Two-dimensional electron gases at a spinel/perovskite complex oxide heterointerface with electron mobilities exceeding $100,000 \text{ cm}^2 \text{V}^{-1} \text{s}^{-1}$

Y. Z. Chen, F. Trier, D. V. Christensen, S. Linderoth, and N. Pryds

Department of Energy Conversion and Storage, Technical University of Denmark, Risø Campus, 4000 Roskilde, Denmark

The high-mobility two-dimensional electron gas (2DEG) confined at the interface of two insulating complex oxides provides opportunities for a new generation of electronic devices. So far, such oxide 2DEG is nearly exclusively created within the frame of interface polarity, such as the case of the intensively explored $\text{LaAlO}_3/\text{SrTiO}_3$ (LAO/STO) heterointerface. Alternatively, when building heterostructures on STO, the basis material for oxide electronics, the conductance can also originate from tunable redox reactions at the interface, i.e. the oxygen-vacancies dominated conductivity in reduced STO substrates [1]. In this presentation, the mechanism of the interface conductance in STO-based oxide heterostructures will be discussed. Moreover, relying on redox reactions, we created a new type of 2DEG at the heterointerface between SrTiO_3 and a spinel $\gamma\text{-Al}_2\text{O}_3$ epitaxial film with compatible oxygen ions sublattices [2]. Electron mobilities more than one order of magnitude higher than those of hitherto investigated perovskite-type interfaces were obtained. The spinel/perovskite 2DEG, where the two-dimensional conduction character is revealed by quantum magnetoresistance oscillations, is found to result from interface-stabilized oxygen vacancies confined within a layer of 0.9 nm in proximity to the interface. Our findings pave the way for studies of mesoscopic physics with complex oxides and design of high-mobility all-oxide electronic devices

1. Y. Z. Chen *et al.*, *Nano Lett.* **11**, 3774 (2011).
2. Y. Z. Chen *et al.*, *Nature Communications*. doi:10.1038/ncomms2394 (2012) (*in progress*).

Corresponding author: Yunzhong Chen, Email: yunc@dtu.dk, Frederiksborgvej 399, DK 4000, Roskilde, Denmark, Tel: +45 4677 5614; Fax: +45 4677 5858